

# MANCOZEB

FUNGICIDE

DISEASE CONTROL



HOW TO USE: (the measuring spoon provided holds 3gms)	DISEASE	RATE per 9L of Water	WHP	HOW TO APPLY
<b>ORNAMENTALS</b>				
Ferns	Botrytis leaf spot	9g	-	Apply when disease symptoms first appear and then repeat at 7-10 day intervals.
Flowers, Shrubs	Septoria leaf spot	18g		Apply when disease symptoms first appear and then repeat at 7-14 day intervals
Carnations	Rust, Alternaria leaf spot			Apply when disease symptoms first appear and then repeat at 7-10 day intervals.
Gladioli	Grey mould (Botrytis)			
Chrysanthemums	Septoria leaf spot, Rust			
Roses	Black Spot			
Turf	Fusarium, Rhizoctonia, Helminthosporium	18g in 9 litres of water per 9 sq metres		Apply 9 litres spray per 9 square metres. Repeat each 7 days during susceptible period. Do not feed treated grass clippings to poultry or animals.
<b>FRUIT</b>				
Apples, Pears	Apple Scab (Black spot), Bitter rot, Fly speck, Pear Scab, Ripe fruit spot, Sooty blotch, Target spot	18g	14 days	Spray at 7-14 day intervals following green tip copper spray. Use shorter spray interval during prolonged wet conditions.
Stone fruit	Brown rot, Freckle, Rust Shot hole			Spray at early bloom, mid to full bloom, petal and shuck fall then at intervals of 14 days.
Mangoes	Anthraxnose	18g		Spray at weekly intervals during flowering then monthly until near harvest.
<b>VEGETABLES</b>				
Beans	Rust, Anthracnose	18g	7 days	Spray at first sign of disease. Repeat at 7-10 day intervals.
Capsicum	Target spot			Spray at first sign of disease. Repeat at 7-10 day intervals.
Cabbage, Cauliflower, Broccoli, Brussel sprouts	Alternaria spot, Anthracnose Downy Mildew, Ring spot			Spray at first sign of disease. Repeat at 7-10 day intervals.
Cucurbit Pumpkins, Melons, Cucumber, Cantaloupe, Zucchini, Squash	Anthraxnose, Downy Mildew, Gummy stem blight, Septoria spot			Spray at first sign of disease. Repeat at 7-10 day intervals while the risk of disease exists.
Lettuce	Anthraxnose, Downy Mildew, Septoria leaf spot		14 days	Spray at first sign of disease. Repeat at 7-10 day intervals. Use shorter intervals during weather conditions favourable to disease development.
Onions & Garlic	Downy mildew, Purple blotch		7 days	
Peas	Downy Mildew			
Potatoes	Early blight (Target spot) Late blight (Irish blight)			Spray at first sign of disease. Repeat at 7-10 day intervals. Do not apply to tomato seedlings before transplanting.
Spinach, Silver Beet, Beetroot	Cercospora leaf spot, Downy mildew			
Tomatoes	Anthraxnose, Early blight, (Target spot), Grey leaf spot, Late blight (Irish blight), Leaf mould, Phoma rot			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED. DO NOT PICK APPLES, MANGOES, STONE FRUIT AND LETTUCE FOR 14 DAYS AFTER SPRAYING. DO NOT PICK VEGETABLES (EXCEPT LETTUCE), MELONS, CANTALOUPE AND TOMATOES FOR 7 DAYS AFTER SPRAYING.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS



DISEASE CONTROL

# MANCOZEB

FUNGICIDE

ACTIVE CONSTITUENT:  
750g/KG MANCOZEB

For the control of  
certain diseases of Apples,  
Mangoes, Vegetables,  
Flowers & Turf.



Net Contents 100g

**HOW TO PREPARE:** Product can be applied through most garden sprayers. Add about half the required volume of water to the sprayer and then add the required amount of product.

The Chemspray measure supplied contains 3 grams.

Mix or agitate and then add the remaining water. Regular agitation during spraying is necessary to avoid product settling out. Do not store mixed product. Only mix what you can use immediately. Thoroughly cover plants with spray ensuring all leaves are wetted. Avoid excessive run-off.

**CAUTION:** DO NOT feed grass clippings from treated areas to poultry or animals. Dangerous to fish. DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

**STORAGE & DISPOSAL:** Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

**SAFETY DIRECTIONS:** May irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. Avoid giving alcohol.

APVMA APPROVAL NO. 30578/100/0305

Batch No: DOM:



**Amgrow Pty Ltd**  
ABN 81 100 684 786  
82 Christensen Road,  
Staplyton QLD 4207  
**Customer Service Number:**  
1800 063 619  
**www.amgrow.com.au**



a 4/106 grose street, parramatta nsw 2150  
p 02 9630 1733 f 02 9630 1930  
e info@dbgraphics.com.au  
w www.dbgraphics.com.au

Name:	Mancozeb 100g carton
DBG Job No:	55658
Item Code:	N/A
Date:	04/05/11
Proof No.:	3
Dimensions:	279 x 265.75 mm (66.5w x 141h x 66.5d mm)
Colours:	C M Y K PMS PANTONE Rhodamine Red PMS 137

**PLEASE EXAMINE THIS PROOF CAREFULLY**

Whilst all care and attention is taken when preparing your design, it is requested you check all details carefully. It is the responsibility of the client to ensure that all details shown are correct and ready for production. The client is responsible for any errors in the design, which are not corrected by the client, before the design is completed. This is a proof, the colours and screen densities used in the proofing process provide a representation only, the final colour(s) may vary from this proof.